

§ 520.447

of *S. aureus*, *B. fragilis*, *P. melaninogenicus*, *F. necrophorum*, and *C. perfringens*.

(ii) For the treatment of soft tissue infections (wounds and abscesses), dental infections, and osteomyelitis caused by susceptible strains of *S. aureus*, soft tissue infections (deep wounds and abscesses), dental infections, and osteomyelitis caused by or associated with susceptible strains of *B. fragilis*, *P. melaninogenicus*, *F. necrophorum*, and *C. perfringens*.

[67 FR 54954, Aug. 27, 2002, as amended at 68 FR 55824, Sept. 29, 2003; 69 FR 32273, June 9, 2004]

§ 520.447 Clindamycin liquid.

(a) *Specifications*. Each milliliter of solution contains the equivalent of 25 milligrams (mg) clindamycin as the hydrochloride salt.

(b) *Sponsors*. See sponsors in § 510.600(c) of this chapter as follows:

(1) Nos. 000009 and 059130 for use as in paragraphs (d)(1)(i)(A), (d)(1)(ii)(A), (d)(2)(i)(A), and (d)(2)(ii)(A) of this section.

(2) No. 059079 for use as in paragraphs (d)(1)(i)(B), (d)(1)(ii)(B), (d)(2)(i)(B), and (d)(2)(ii)(B) of this section.

(c) *Special considerations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d) *Conditions of use*—(1) *Dogs*—(i) *Amount*—(A) Wounds, abscesses, and dental infections: 2.5 to 15 mg per pound (lb) of body weight every 12 hours for a maximum of 28 days. Osteomyelitis: 5.0 to 15 mg/lb of body weight every 12 hours for a minimum of 28 days.

(B) Wounds, abscesses, and dental infections: 2.5 mg per pound (lb) of body weight every 12 hours for a maximum of 28 days. Osteomyelitis: 5.0 mg/lb of body weight every 12 hours for a minimum of 28 days.

(ii) *Indications for use*—(A) For the treatment of skin infections (wounds and abscesses) due to susceptible strains of coagulase-positive staphylococci (*Staphylococcus aureus* or *S. intermedius*), deep wounds and abscesses due to susceptible strains of *Bacteroides fragilis*, *Prevotella melaninogenicus*, *Fusobacterium necrophorum*, and *Clostridium perfringens*, dental infections due to

21 CFR Ch. I (4–1–05 Edition)

susceptible strains of *S. aureus*, *B. fragilis*, *P. melaninogenicus*, *F. necrophorum*, and *C. perfringens*, and osteomyelitis due to susceptible strains of *S. aureus*, *B. fragilis*, *P. melaninogenicus*, *F. necrophorum*, and *C. perfringens*.

(B) For the treatment of soft tissue infections (wounds and abscesses), dental infections, and osteomyelitis caused by susceptible strains of *S. aureus* and for soft tissue infections (deep wounds and abscesses), dental infections, and osteomyelitis caused by or associated with susceptible strains of *B. fragilis*, *P. melaninogenicus*, *F. necrophorum*, and *C. perfringens*.

(2) *Cats*—(i) *Amount*—(A) 5.0 to 15.0 mg/lb of body weight every 24 hours for a maximum of 14 days.

(B) 5.0 to 10.0 mg/lb of body weight every 24 hours for a maximum of 14 days.

(ii) *Indications for use*—(A) For the treatment of skin infections (wounds and abscesses) due to susceptible strains of *S. aureus*, *S. intermedius*, *Streptococcus* spp., deep wounds and abscesses due to susceptible strains of *Clostridium perfringens* and *Bacteroides fragilis*, and dental infections due to susceptible strains of *S. aureus*, *S. intermedius*, *Streptococcus* spp., *C. perfringens*, and *B. fragilis*.

(B) *Aerobic bacteria*: Treatment of soft tissue infections (wounds and abscesses) and dental infections caused by or associated with susceptible strains of *S. aureus*, *S. intermedius*, and *Streptococcus* spp. *Anaerobic bacteria*: Treatment of soft tissue infections (deep wounds and abscesses) and dental infections caused by or associated with susceptible strains of *C. perfringens* and *B. fragilis*.

[67 FR 54954, Aug. 27, 2002, as amended at 67 FR 78684, Dec. 26, 2002; 68 FR 55824, Sept. 29, 2003; 69 FR 31734, June 7, 2004]

§ 520.452 Clenbuterol syrup.

(a) *Specifications*. Each milliliter contains 72.5 micrograms of clenbuterol hydrochloride.

(b) *Sponsor*. See 000010 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Horses*—(i) *Amount*. Administer orally twice a day (b.i.d.). Initial dose is 0.5 milliliter per